

PAPILIO AMPHRYSIUS.



CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes ;
(volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis denticulatis nigris, versus apices flavo-substriatis, inferioribus flavis nigro-marginatis.

PAPILIO alis concoloribus nigris, anterioribus maculis, posterioribus disco flavis.

Lin. Syst. Nat. Gmel. p. 2230.

Eq. Tr.


Merian Ins. Surin. t. 72 ♀

PAPILIO AMPHRYSIUS.

Cram. t. 219. f. A.

Affinis P. Helenæ.

Insulas Javam atque Amboinam præcipue incolit formosa hæc papilionis species, magnitudine vera in tabula depicta.





R.P. Neder. D. el: a. Sulp

AMPHRYSIUS.



GENERIC CHARACTER.

Antennæ generally thickening towards the extremities, and terminating in a elavated tip.
Wings (when sitting) erect, and meeting upwards.

(Flight diurnal.)

SPECIFIC CHARACTER, &c.

BUTTERFLY with denticulated wings; the upper pair blaek, with obscure yellow streaks towards the tips; the lower yellow, bordered with blaek.

Jablonsk. pap. 1. t. 1. f. 3.

This beautiful Butterfly is principally found in the islands of Java and Amboina, and is represented in its natural size on the annexed plate.